

# High $p_T$ Results from

Thomas Peitzmann



Utrecht University/NIKHEF



for the STAR collaboration

# Outline


- High- $p_T$  Inclusive Spectra
- High- $p_T$  Dihadron Correlations
- Intermediate- $p_T$  Near-Side
- Intermediate- $p_T$  Away-Side
- Heavy Flavour
- The Future: Photons

# High- $p_T$ Inclusive Spectra

Where's the fragmentation regime?  
What's the lower limit on the medium density?


# Inclusive Suppression

J. Adams et al, Phys. Rev. Lett. 91 (2003) 072304




- studied with nuclear modification factor
 
$$R_{AA}(p_T) = \frac{d^2N^{AA} / dp_T d\eta}{T_{AA} d^2\sigma^{NN} / dp_T d\eta}$$
- established probe of final state suppression
  - sensitive to density of the medium
- very close to maximum suppression
  - provides only lower bound on density
- surface bias in high  $p_T$  hadron emission!

# Dependence of Suppression on Geometry




- Cu+Cu fit smoothly to trend of Au+Au
  - smaller uncertainties for small  $N_{\text{part}}$
- universal dependence on  $N_{\text{part}}$ 
  - does not strongly favour particular scaling law

# Baryon vs. Meson $R_{cp}$



- baryon/meson separation at intermediate  $p_T$ 
  - baryon enhancement
- explained by non-fragmentation contribution
  - e.g. recombination
- disappears at  $p_T \approx 6$  GeV/c
  - fragmentation limit?
  - difference between q and g energy loss?

# $v_2$ of Identified Hadrons



- meson/baryon scaling extends out to high  $p_T$
- possibly: surface bias yields identical  $R_{AA}$  ( $R_{cp}$ ) in central collisions for  $q$  and  $g$ , but difference still remains in  $v_2$  from non-central collisions

PHENIX (open symbols):  
Phys. Rev. Lett. 91, 182301 (2003)


T. Peitzmann, Hard Probes 2006, Asilomar

- finite  $v_2$  at high  $p_T$
- saturation and onset of decline at  $p_T > 3$  GeV/c
- clear meson/baryon scaling at intermediate  $p_T$
- $v_2$  from anisotropy in jet quenching
  - path length dependence

# High- $p_T$ Dihadron Correlations

Can we get an upper limit on the medium density?  
How does fragmentation work after energy loss?


# Dihadrons in Heavy Ion Collisions



- near-side correlation unchanged
- away-side peak suppressed in central  $Au+Au$ 
  - consistent with surface emission
- distributions in  $d\text{Au}$  similar to  $p+p$ 
  - suppression is final state effect
- **more intuitive hint for “jet” suppression, but quantitatively much more difficult**


# Emergence of Di-Jets

$8 < p_T(\text{trig}) < 15 \text{ GeV}/c$




- clear away side peak even in central Au+Au for high trigger  $p_T$
- background reduced for higher associated  $p_T$
- little modification of near-side yield
- suppression of away-side yield apparent in central Au+Au


# Width of Away-Side Peaks



- away-side widths similar for central and peripheral
- fragmentation as in vacuum?



# Momentum Distribution of Near-Side Yield




- Modified di-hadron fragmentation function (X.-N. Wang)

$$D^{h_1 h_2}(z_T, p_T^{\text{trig}}) = p_T^{\text{trig}} \frac{d\sigma_{AA}^{h_1 h_2} / dp_T^{\text{trig}} dp_T}{d\sigma_{AA}^{h_1} / dp_T^{\text{trig}}}$$


$$z_T \equiv \frac{p_T^{\text{assoc}}}{p_T^{\text{trig}}}$$

- near-side yield consistent with no modification
  - no dependence of ratio on  $z_T$  in measured range

# Momentum Distribution of Away-Side Yield




- away-side yield strongly suppressed
  - level of  $R_{AA}$
  - different surface bias?
  - upper limit on medium density obtainable?
- no dependence of ratio on  $z_T$  in measured range
  - vacuum fragmentation?



# Intermediate- $p_T$ Near-Side



Do we see the coupling of the jet to the medium?  
Can we test recombination models?

# Near-Side Long-Range $\Delta\eta$ Correlation: the Ridge



# Centrality Dependence of the Ridge


$3 < p_{t,\text{trigger}} < 4 \text{ GeV}$  and  $p_{t,\text{assoc.}} > 2 \text{ GeV}$



J. Putschke et al.,  
parallel talk


- yield of associated particles can be separated into a jet-like yield and a ridge yield
  - jet-like yield consistent in  $\eta$  and  $\phi$  and independent of centrality
  - ridge yield increases with centrality

# Rapidity Dependence



- approximately linear dependence on rapidity
  - longitudinal scaling
    - relation to longitudinal flow?
  - shorter range for larger trigger  $p_T$


# Dependence on Trigger $P_T$



- yield significant up to  $P_T = 9 \text{ GeV}/c$  in central
  - clearly jet-related
- no strong dependence on trigger hadron momentum
  - increase of jet-like yield  $\Rightarrow$  decrease of relative yield

# Shape of Associated Particle Spectra

$$dN/dp_t \propto p_t e^{-p_t/T}$$





- jet-like spectra harder than inclusive
  - flatter for higher trigger  $p_T$
- ridge spectra similar to inclusive
  - slightly larger slope
  - approximately independent of trigger  $p_T$



# Recombination and Dihadron Correlations


- large baryon yield at intermediate  $p_T$  explained by recombination models
- difference in correlation structure expected
  - naïve expectation from pure thermal reco: no correlation
  - more realistic models: some correlation due to thermal-shower reco (R. Hwa)
- hadrons from recombination should show a modified correlation
  - larger fraction of baryons produced from recombination (compared to mesons)
  - stronger modification of correlation structure expected for baryon triggers

# Identified Hadron Triggers



- total associated yield increases with centrality: **ridge!**
- jet-like yield independent of centrality
  - differences for strange particles

# Identified Hadron Triggers



- less associated jet-like yield for  $\Lambda$ 
  - higher ridge yield for  $\Lambda$
  - consistent with recombination?
- attention when extracting correlation strength in  $\Delta\phi$  only!


# Interpretation of the Ridge

- coupling of high energy parton to longitudinal flow (Armesto et al, nucl-ex/0405301)
  - expect broadening but not plateau
- correlation from radial flow (Voloshin nucl-th/0312065)
  - not expected at high  $P_T$
- thermal recombination + local heating from energy loss (Chiu & Hwa, Phys. Rev. C72 034903, 2005)
  - qualitatively consistent
- more general:
  - ridge is jet-related structure with properties similar to bulk
  - early coupling: before longitudinal expansion
  - local energy (heating) or momentum transfer (collective flow)
- Does ridge measure the amount of energy transferred to the bulk?

# Intermediate- $p_T$ Away-Side




Where does the jet energy go?  
Is their conical flow?

# Two-Particle Correlations (Mach Cone?)




- broad away-side distribution in central Au+Au
  - enhanced yield for lower  $p_T$
  - consistent with two-peak structure
    - Mach cone or deflected jets?  
⇒ study 3-part. correlation
    - sensitive to elliptic flow subtraction
- dependence on trigger  $p_T$ ?
- enhanced yield for near-side
  - quantitatively consistent with ridge
  - **near-side enhancement only ridge?**  
⇒ vacuum fragmentation?



# Conical Flow vs Deflected Jets




# Three-Particle Correlations



Raw

Jet x Bkgd  
(Hard-Soft)Bkgd x Bkgd  
(Soft-Soft incl. Flow)

Au+Au Central 0-12% Triggered




- signal obtained by subtraction of dominant backgrounds
  - flow components, jet-related two-particle correlation
- improved analysis compared to QM (e.g. high statistics)
  - additional check with cumulant analysis under way
  - careful: different assumptions on background normalisation!
- clear elongation (jet deflection)
- off-diagonal signal related to mach cone?

# Heavy Flavour


Do we understand heavy flavour production?  
What is the energy loss mechanism?

# Electrons in p+p Compared to pQCD



- single (non-photonic) electrons mainly from c and b
- pQCD (FONLL) scaled by K-factor  $\approx 5.7$  to match the data
- Ratio Data/pQCD is independent of  $p_T$  for  $p_T < 8$  GeV/c
  - same K-factor for charm and beauty?


# Single Electron $R_{AA}$



Armesto et al.  
van Hees et al.  
Wicks et al. (DVGL)

hep-th/0511257  
Phys. Rev. C 73, 034913 (2006)  
hep-th/0512076

## Charged Hadron $R_{AA}$



theoretical description needs:

- extremely high density or
- significant contributions of collisional energy loss and dominance of charm up to high  $p_T$

- $R_{AA}$  to 10 GeV/c in non-photonic electrons
- suppression is approximately the same as for light hadrons



# The Future: Photons

The reference probe.

Can we tag the initial jet energy?

Can we measure energy loss?


d+Au 2003



T. Peitzmann, Hard Probes 2006, Asilomar

- measured with STAR Barrel EMC
- reconstruction of two-photon invariant mass
- agreement with NLO pQCD in d+Au
- used as input for direct photon analysis
  - statistical subtraction of decay background


# Direct Photons in d+Au



$$R \equiv \frac{\left(\gamma_{\text{incl}}/\pi^0\right)_{\text{measured}}}{\left(\gamma_{\text{decay}}/\pi^0\right)_{\text{simulated}}} \approx 1 + \frac{\gamma_{\text{direct}}}{\gamma_{\text{decay}}}$$


- proof of principle measurement with STAR Barrel EMC
- setting a baseline for Au+Au
- consistent with NLO pQCD
- reduction of systematic errors needed to extract spectrum

# Towards $\gamma$ -Jet in Au+Au



- photon tag
  - potentially clean jet preparation
- clear near-side and away-side correlation peaks
- strong contamination from  $\pi^0$  decay photons
  - reduction in near-side strength compatible with direct photon component
- work in progress ...

# Shower Shape Analysis in d+Au



- charged hadrons associated with high  $E_T$  EM shower
- shower width selects  $\pi^0$  and  $\gamma$  candidates
- measure associated yield for  $\pi^0$  triggers
  - reduction of yield can be used to estimate direct  $\gamma$  contribution in other samples
- possibility to enrich trigger sample with direct  $\gamma$ !

# Summary

- fragmentation regime reached at  $p_T = 6 \text{ GeV}/c$ 
  - elliptic flow consistent?
- upper limit on density from di-hadron correlations?
- jet-like yield consistent with vacuum fragmentation
  - width, yield and momentum spectrum unchanged
- enhancement on near-side due to soft correlation
  - long range in  $\eta$  (ridge)
  - ridge yield indicates amount of energy loss?
  - careful in  $\Delta\phi$ -only analysis!
- softening and broadening on the away-side
  - 2-part. correlations consistent with Mach cone
  - 3-part. correlations show jet-deflection - Mach cone signal not well established
- heavy flavour
  - importance of c and b
  - energy loss mechanism?
- photons
  - direct photons in d+Au
  - getting ready to exploit EMC capabilities for photon tagged jets



# The STAR Collaboration

## U.S. Labs:

Argonne, Lawrence Berkeley, and Brookhaven National Labs

## U.S. Universities:

UC Berkeley, UC Davis, UCLA, Caltech, Carnegie Mellon, Creighton, Indiana, Kent State, MIT, MSU, CCNY, Ohio State, Penn State, Purdue, Rice, Texas A&M, UT Austin, Washington, Wayne State, Valparaiso, Yale

## Brazil:

Universidade de Sao Paolo

## China:

IHEP - Beijing, IPP - Wuhan, USTC, Tsinghua, SINAP, IMP Lanzhou

## Croatia:

Zagreb University

## Czech Republic:

Nuclear Physics Institute

## England:

University of Birmingham

## France:

Institut de Recherches Subatomiques Strasbourg, SUBATECH - Nantes

## Germany:

Max Planck Institute – Munich University of Frankfurt

## India:

Bhubaneswar, Jammu, IIT-Mumbai, Panjab, Rajasthan, VECC

## Netherlands:

NIKHEF/Utrecht

## Poland:

Warsaw University of Technology

## Russia:

MEPHI – Moscow, LPP/LHE JINR – Dubna, IHEP – Protvino

## South Korea:

Pusan National University

## Switzerland:

University of Bern